



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

INDEX.

VAN VLECK, E. B., On the Determination of a Series of Sturm's Functions by the Calculation of a Single Determinant, . . .	1
WHITE, H. S., Two Elementary Geometrical Applications of Determinants,	103
WHITTEMORE, J. K., Note on the Convergence of Definite Integrals, . . .	189
WILSON, E. B., Note on the Function Satisfying the Functional Relation $\phi(u) \cdot \phi(v) = \phi(u + v)$,	47

ERRATA.

Page 21. In the figure θ and ϕ should be interchanged.

Page 81. The following foot note was omitted: This paper was read before the American Mathematical Society at the Meeting of August 25, 1899.

Page 111, line 11 The equation should read: $y - y' = 8x'^2(x - x')$.

Page 114, line 10. For $+6x^2$ read $-6x^2$.

Page 115, line 9. For $R =$ read $R' =$.

Page 119, line 1. For c' read b' .